

IMMUNE MEDIATED SKIN INFLAMMATION IS THE SAME IN ATOPIC DERMATITIS PATIENTS WITH OR WITHOUT FILAGGRIN MUTATION

Mócsai Gábor^{1,2}

Dajnoki Zsolt^{1,2}, Kapitány Anikó^{1,2}, Béke Gabriella^{1,2}, Gáspár Krisztián^{1,2}, Emri Gabriella², Nagy Bence³, Dezső Balázs³, Szegedi Andrea^{1,2}

¹Division of Dermatological Allergology, ²Department of Dermatology, ³Department of Pathology, Faculty of Medicine, University of Debrecen, Debrecen, Hungary

Introduction: Recent investigations showed that T helper (Th) 2 inflammatory cytokines can impair skin barrier. The opposite question, whether skin barrier alterations may affect keratinocyte (KC) immune responses, remained less studied. Our aim was to investigate whether the immune mediated skin inflammation, KC function, T and dendritic cell (DC) count differ between severe atopic dermatitis (AD) patients with or without filaggrin (FLG) mutation.

Methods: Lesional skin biopsies of severe FLG mutant and wild type (Wt) AD patients and controls were analyzed by immunohistochemistry: FLG and inflammatory T(h)2 polarizing cytokines (TSLP, IL-33) and CCL27 chemokine, characteristically expressed by KCs in lesional AD skin, were immunostained. Histological severity markers (Ki67 positivity and epidermal thickness) as well as CD3⁺ T cell and CD11c⁺ DC count were also investigated. Whole slide imaging method allowed us to compare protein levels. Correlations between cytokines/chemokine and severity markers were calculated.

Results: Compared to controls, in both AD groups, FLG levels were found significantly lower, whilst the same level of FLG expression was measured in the AD groups. Concerning the levels of the investigated parameters no significant differences were found between the two AD groups. The expression of AD specific cytokines and chemokine showed significant correlation with histological severity markers.

Conclusions: Our findings suggest that the immune mediated skin inflammation in FLG mutant and Wt AD patients is mediated by the same amount of immune cells and KC derived cytokines and chemokines. Results of the correlations demonstrated that KC's immune activation is connected rather to the severity of the disease, then to the origin of barrier alterations.

Poszter

Klinikai